

### **VASODIP®**

(Lercanidipine hydrochloride)

Film-coated tablets

### **QUALITATIVE AND QUANTITATIVE COMPOSITION:**

One tablet of Vasodip 10 contains as active ingredient: 10 mg of lercanidipine hydrochloride, which is equivalent to 9.4 mg of lercanidipine.

One tablet of Vasodip 20 contains as active ingredient: 20 mg of lercanidipine hydrochloride, which is equivalent to 18.8 mg of lercanidipine.

For excipients, see Pharmaceutical particulars.

### **CLINICAL PARTICULARS**

#### **Therapeutic indications:**

Vasodip is indicated for the treatment of mild to moderate essential hypertension. Some individuals not adequately controlled on a single antihypertensive agent, may benefit from the addition of Vasodip to therapy with a beta-adrenoceptor blocking drug (atenolol), a diuretic (hydrochlorothiazide) or an angiotensin converting enzyme inhibitor (captopril or enalapril).

#### **Posology and method of administration:**

The recommended dosage is 10 mg orally once a day at least 15 minutes before meals, the dose may be increased to 20 mg depending on the individual patient's response.

Dose titration should be gradual, because it may take about 2 weeks before the maximal antihypertensive effect is apparent.

Since the dose-response curve is steep with a plateau at doses between 20-30 mg, it is unlikely that efficacy will be improved by higher doses; whereas side effects may increase.

*Use in the elderly:* Although the pharmacokinetic data and clinical experience suggest that no adjustment of the daily dosage is required, special care should be exercised when initiating treatment in the elderly.

*Use in children:* Since there is no clinical experience in patients under the age of 18 years, use in children is not currently recommended.

*Use in renal or hepatic dysfunction:* special care should be exercised when treatment is commenced in patients with mild to moderate renal or hepatic dysfunction. Although the usually recommended dose schedule may be tolerated by these subgroups, an increase in dose to 20 mg daily must be approached with caution.

The antihypertensive effect may be enhanced in patients with hepatic impairment and consequently an adjustment of the dosage should be considered. Vasodip is not recommended for use in patients with severe hepatic impairment or in patients with severe renal impairment (GFR < 30 ml/min).

### **Contraindications:**

- Hypersensitivity to the active substance “lercanidipine“, to any dihydropyridine or to any of the excipients of the medicinal product.
- Pregnancy and lactation (see: Pregnancy and lactation).
- Women of child-bearing potential unless effective contraception is used.
- Left ventricular outflow tract obstruction.
- Untreated congestive cardiac failure.
- Unstable angina pectoris.
- Severe renal or hepatic impairment.
- Within 1 month of a myocardial infarction.
- Co-administration with:
  - Strong inhibitors of CYP3A4.
  - Cyclosporin
  - Grapefruit juice.(see: Interaction with other medicinal products and other forms of interaction).

### **Special warnings and precautions for use:**

Special care should be exercised when Vasodip is used in patients with sick sinus syndrome (if a pacemaker is not in situ). Although hemodynamic controlled studies revealed no impairment of ventricular function, care is also required in patients with LV dysfunction. It has been suggested that some short-acting dihydropyridines may be associated with increased cardiovascular risk in patients with ischaemic heart disease. Although Vasodip is long acting, caution is required in such patients. Some dihydropyridines may rarely lead to precordial pain or angina pectoris. Very rarely patients with pre-existing angina pectoris may experience increased frequency, duration or severity of these attacks. Isolated cases of myocardial infarction may be observed (see: Undesirable effects).

*Use in renal or hepatic dysfunction:* special care should be exercised when treatment is commenced in patients with mild to moderate renal or hepatic dysfunction. Although the usually recommended dose schedule may be tolerated by these subgroups, an increase in dose to 20 mg daily must be approached with caution. The antihypertensive effect may be enhanced in patients with hepatic impairment and consequently an adjustment of the dosage should be considered. Vasodip is not recommended for use in patients with severe hepatic impairment or in patients with severe renal impairment (GFR < 30 ml/min) (see: Posology and method of administration).

Alcohol should be avoided since it may potentiate the effect of vasodilating antihypertensive drugs (see: Interaction with other medicinal products and other forms of interaction).

Inducers of CYP3A4 like anticonvulsants (e.g. phenytoin, carbamazepine) and

rifampicin may reduce lercanidipine's plasma levels and therefore the efficacy of lercanidipine may be less than expected. (see: Interaction with other medicinal products and other forms of interaction).

One tablet of Vasodip 10 contains 30 mg lactose. This amount is probably too small to give rise to significant symptoms in patients with lactose intolerance. 1 tablet of Vasodip 20 contains 60 mg lactose and therefore should not be administered to patients with Lapp lactase insufficiency, galactosaemia or glucose/galactose malabsorption syndrome.

### **Interaction with other medicinal products and other forms of interaction:**

Lercanidipine is known to be metabolised by the CYP3A4 enzyme and therefore inhibitors and inducers of CYP3A4 administered concurrently may interact with the metabolism and elimination of lercanidipine.

Co-prescription of Vasodip with inhibitors of CYP3A4 (e.g. ketoconazole, itraconazole, ritonavir, erythromycin, troleandomycin) should be avoided (see: Contraindications). An interaction study with a strong CYP3A4 inhibitor, Ketoconazole, has shown a considerable increase in plasma levels of lercanidipine (a 15-fold increase of the AUC and an 8-fold increase of the  $C_{max}$  for the eutomer S-lercanidipine).

Cyclosporin and lercanidipine should not be administered together (see: Contraindications). Increased plasma levels of both lercanidipine and cyclosporin have been observed following concomitant administration. A study in young healthy volunteers has shown that when cyclosporin was administered 3 hours after the lercanidipine intake the plasma levels of lercanidipine did not change, while the AUC of cyclosporin increased by 27%. However, the co-administration of Vasodip with cyclosporin has caused a 3-fold increase of the plasma levels of lercanidipine and a 21% increase of the cyclosporin AUC.

Lercanidipine should not be taken with grapefruit juice (see: Contraindications). As for other dihydropyridines, lercanidipine is sensitive to inhibition of metabolism by grapefruit juice, with a consequent rise in its systemic availability and increased hypotensive effect.

When concomitantly administered at a dose of 20 mg with midazolam p.o. to elderly volunteers, lercanidipine's absorption was increased (by approximately 40%) and the rate of absorption was decreased ( $t_{max}$  was delayed from 1.75 to 3 hours). Midazolam concentrations were not modified.

Caution should be exercised when Vasodip is co-prescribed with other substrates of CYP 3A4, like terfenadine, astemizole, class III antiarrhythmic drugs such as amiodarone and quinidine.

Co-administration of Vasodip with CYP 3A4 inducers like anticonvulsants (e.g. phenytoin, carbamazepine) and rifampicin should be approached with caution since the antihypertensive effect may be reduced and blood pressure should be monitored more frequently than usual.

When Vasodip was co-administered with metoprolol, a  $\beta$ -blocker eliminated mainly by the liver, the bioavailability of metoprolol was not changed while that of lercanidipine was reduced by 50%. This effect may be due to the reduction in the

hepatic blood flow caused by  $\beta$ -blockers and may therefore occur with other drugs of this class. Consequently, lercanidipine may be safely administered with beta-adrenoceptor blocking drugs, but dose adjustment may be required. An interaction study with fluoxetine (an inhibitor of CYP2D6 and CYP3A4), conducted in volunteers of an age of  $65 \pm 7$  years (mean  $\pm$  s.d.), has shown no clinically relevant modification of the pharmacokinetics of lercanidipine.

Concomitant administration of cimetidine 800 mg daily does not cause significant modifications in plasma levels of lercanidipine, but at higher doses caution is required since the bioavailability and the hypotensive effect of lercanidipine may be increased.

Co-administration of 20 mg lercanidipine in patients chronically treated with  $\beta$ -methyl digoxin showed no evidence of pharmacokinetic interaction. Healthy volunteers treated with digoxin following dosing with 20 mg lercanidipine given fasted, showed a mean increase of 33% in digoxin  $C_{max}$  while AUC and renal clearance were not significantly modified. Patients on concomitant digoxin treatment should be closely monitored clinically for signs of digoxin toxicity.

When a dose of 20 mg of Vasodip was repeatedly co-administered with 40 mg of simvastatin, the AUC of lercanidipine was not significantly modified, while simvastatin's AUC increased by 56% and that of its active metabolite  $\beta$ -hydroxyacid by 28%. It is unlikely that such changes are of clinical relevance. No interaction is expected when lercanidipine is administered in the morning and simvastatin in the evening, as indicated for such drug.

The co-administration of 20 mg lercanidipine to healthy volunteers given fasted did not alter the pharmacokinetics of warfarin.

Vasodip has been safely administered with diuretics and ACE inhibitors.

Alcohol should be avoided since it may potentiate the effect of vasodilating antihypertensive drugs (see: Special warnings and precautions for use).

### **Pregnancy and lactation:**

Data for lercanidipine provide no evidence of a teratogenic effect in the rat and the rabbit and reproductive performance in the rat was unimpaired. Nevertheless, since there is no clinical experience with lercanidipine in pregnancy and lactation, and other dihydropyridine compounds have been found teratogenic in animals, Vasodip should not be administered during pregnancy or to women with child-bearing potential unless effective contraception is used. Because of high lipophilicity of lercanidipine, distribution in milk may be expected. Therefore, it should not be administered to nursing mothers.

### **Effects on ability to drive and use machines:**

Clinical experience with lercanidipine indicates that it is unlikely to impair a patient's ability to drive or use machinery. However, caution should be exercised because dizziness, asthenia, fatigue and rarely somnolence may occur.

### **Undesirable effects:**

About 1.8% of treated patients experienced adverse reactions.

The table below shows the incidence of adverse drug reactions, at least possibly causally related, grouped by WHO-ART Body System classification, and ranked by frequency (uncommon, rare).

As shown in the table, the most commonly occurring adverse drug reactions reported in controlled clinical trials are headache, dizziness, peripheral oedema, tachycardia, palpitations, flushing, each occurring in less than 1% of patients.

Skin And Appendages Disorders	Rare (>1/10,000 <1/1000)	rash
Musculo-Skeletal System Disorders	Rare (>1/10,000 <1/1000)	myalgia
Central & Peripheral Nervous System Disorders	Uncommon (>1/1000 <1/100)	headache; dizziness
Psychiatric Disorders	Rare (>1/10,000 <1/1000)	somnolence
Gastro-Intestinal System Disorders	Rare (>1/10,000 <1/1000)	nausea; dyspepsia; diarrhoea; abdominal pain; vomiting
Cardiovascular Disorders, General	Uncommon (>1/1000 <1/100)	oedema peripheral
Myo-, Endo- Pericardial & Valve Disorders	Rare (>1/10,000 <1/1000)	angina pectoris
Heart Rate and Rhythm Disorders	Uncommon (>1/1000 <1/100)	tachycardia; palpitations
Vascular (Extracardiac) Disorders	Uncommon (>1/1000 <1/100)	flushing
Urinary System Disorders	Rare (>1/10,000 <1/1000)	polyuria
Body As A Whole - General Disorders	Rare (>1/10,000 <1/1000)	asthenia; fatigue

In post-marketing experience, from spontaneous reports the following undesirable effects were reported very rarely (<1/10,000): gingival hypertrophy, reversible increases in serum levels of hepatic transaminases, hypotension, urinary frequency and chest pain.

Some dihydropyridines may rarely lead to precordial pain or angina pectoris. Very rarely patients with pre-existing angina pectoris may experience increased frequency, duration or severity of these attacks. Isolated cases of myocardial infarction may be observed.

Lercanidipine does not appear to influence adversely blood sugar or serum lipid levels.

### **Overdose:**

In the post-marketing experience, two cases of overdose were reported (150 mg

and 280 mg of lercanidipine, respectively, ingested in an attempt to commit suicide). The first patient developed sleepiness and was treated by gastric lavage. The second patient developed cardiogenic shock with severe myocardial ischaemia and mild renal failure and was treated with high-dose catecholamines, furosemide, digitalis and parenteral plasma expanders. Both cases resolved without sequelae. As with other dihydropyridines, overdosage might be expected to cause excessive peripheral vasodilatation with marked hypotension and reflex tachycardia. In case of severe hypotension, bradycardia and unconsciousness, cardiovascular support could be helpful, with intravenous atropine for bradycardia.

In view of the prolonged pharmacological effect of lercanidipine, it is essential that the cardiovascular status of patients who take an overdose is monitored for 24 hours at least. There is no information on the value of dialysis. Since the drug is highly lipophilic, it is most probable that plasma levels are no guide to the duration of the period of risk and dialysis may not be effective.

## **PHARMACOLOGICAL PROPERTIES**

### **Pharmacodynamic properties:**

Lercanidipine is a calcium antagonist of the dihydropyridine group and inhibits the transmembrane influx of calcium into cardiac and smooth muscle. The mechanism of its antihypertensive action is due to a direct relaxant effect on vascular smooth muscle thus lowering total peripheral resistance. Despite its short pharmacokinetic plasma half-life, lercanidipine is endowed with a prolonged antihypertensive activity because of its high membrane partition coefficient, and is devoid of negative inotropic effects due to its high vascular selectivity. Since the vasodilatation induced by Vasodip is gradual in onset, acute hypotension with reflex tachycardia has rarely been observed in hypertensive patients.

As for other asymmetric 1,4-dihydropyridines, the antihypertensive activity of lercanidipine is mainly due to its (S)-enantiomer.

In addition to the clinical studies conducted to support the therapeutic indications, a further small uncontrolled but randomised study of patients with severe hypertension (mean  $\pm$  SD diastolic blood pressure of  $114.5 \pm 3.7$  mmHg) showed that blood pressure was normalised in 40% of the 25 patients on 20 mg once daily dose and in 56% of 25 patients on 10 mg twice daily doses of Vasodip. In a double-blind, randomized, controlled study versus placebo in patients with isolated systolic hypertension Vasodip was efficacious in lowering systolic blood pressure from mean initial values of  $172.6 \pm 5.6$  mmHg to  $140.2 \pm 8.7$  mmHg.

### **Pharmacokinetic properties:**

Vasodip is completely absorbed after 10-20 mg oral administration and peak plasma levels,  $3.30$  ng/ml  $\pm$   $2.09$  s.d. and  $7.66$  ng/ml  $\pm$   $5.90$  s.d. respectively, occur about 1.5-3 hours after dosing.

The two enantiomers of lercanidipine show a similar plasma level profile: the time to peak plasma concentration is the same, the peak plasma concentration and AUC are, on average, 1.2-fold higher for the (S) enantiomer and the elimination half-lives of the two enantiomers are essentially the same. No "in vivo"

interconversion of enantiomers is observed.

Due to the high first pass metabolism the absolute bioavailability of Vasodip orally administered to patients under fed conditions is around 10%, although it is reduced to 1/3 when administered to healthy volunteers under fasting conditions. Oral availability of lercanidipine increases 4 fold when Vasodip is ingested up to 2 hours after a high fat meal. Accordingly, Vasodip should be taken before meals. Distribution from plasma to tissues and organs is rapid and extensive. The degree of serum protein binding of lercanidipine exceeds 98%. Since plasma protein levels are reduced in patients with severe renal or hepatic dysfunction, the free fraction of the drug may be increased.

Vasodip is extensively metabolised by CYP3A4, no parent drug is found in the urine or the faeces. It is predominantly converted to inactive metabolites and about 50% of the dose is excreted in the urine.

"In vitro" experiments with human liver microsomes have demonstrated that lercanidipine shows some degree of inhibition of CYP3A4 and CYP2D6, at concentrations 160- and 40-fold, respectively, higher than those reached at peak in the plasma after the dose of 20 mg.

Moreover, interaction studies in humans have shown that lercanidipine did not modify the plasma levels of midazolam, a typical substrate of CYP3A4, or of metoprolol, a typical substrate of CYP2D6. Therefore, inhibition of biotransformation of drugs metabolised by CYP3A4 and CYP2D6 by Vasodip is not expected at therapeutic doses.

Elimination occurs essentially by biotransformation.

A mean terminal elimination half life of 8-10 hours was calculated and the therapeutical activity lasts for 24 hours because of its high binding to lipid membrane. No accumulation was seen upon repeated administration.

Oral administration of Vasodip leads to plasma levels of lercanidipine not directly proportional to dosage (non-linear kinetics). After 10, 20 or 40 mg, peak plasma concentrations observed were in the ratio 1:3:8 and areas under plasma concentration time curves in the ratio 1:4:18, suggesting a progressive saturation of first pass metabolism. Accordingly, availability increases with dosage elevation. In elderly patients and in patients with mild to moderate renal dysfunction or mild to moderate hepatic impairment the pharmacokinetic behaviour of lercanidipine was shown to be similar to that observed in the general patient population, patients with severe renal dysfunction or dialysis-dependent patients showed higher levels (about 70%) of the drug. In patients with moderate to severe hepatic impairment, the systemic bioavailability of lercanidipine is likely to be increased since the drug is normally metabolised extensively in the liver.

### **Pre-clinical safety data:**

Safety pharmacological studies in animals have shown no effects on the autonomic nervous system, the central nervous system or on gastrointestinal function at antihypertensive doses.

The relevant effects which have been observed in long-term studies in rats and dogs were related, directly or indirectly, to the known effects of high doses of Ca-

antagonists, predominantly reflecting exaggerated pharmacodynamic activity. Lercanidipine was not genotoxic and showed no evidence of carcinogenic hazard. Fertility and general reproductive performance in rats were unaffected by treatment with lercanidipine.

There was no evidence of any teratogenic effect in rats and rabbits, however, in rats, lercanidipine at high dose levels induced pre-and post implantation losses and delay in foetal development.

Lercanidipine hydrochloride, when administered at high dose (12 mg/kg/day) during labour, induced dystocia.

The distribution of lercanidipine and/or its metabolites in pregnant animals and their excretion in breast milk have not been investigated.

Metabolites have not been evaluated separately in toxicity studies.

## **PHARMACEUTICAL PARTICULARS:**

### **List of excipients:**

The tablet core excipients are: lactose, microcrystalline cellulose, sodium starch glycolate, povidone, magnesium stearate.

The film coating contains hypromellose, talc, titanium dioxide, macrogol 6000, ferric oxide.

### **How supplied:**

Film-coated tablets in blister packs.

Store in a dry place at room temperature.

The format of this leaflet was determined by the Ministry of Health (MOH) and its content was checked and approved by the MOH.



## **MANUFACTURER**

**DEXCEL LTD.** A **DEXON** COMPANY

Or-Akiva, P.O.B. 50, Hadera 38100, Israel, Tel: 04-6364000, Fax: 04-6364004.